Observational entropy, coarse-grained states, and irretrodictability

Francesco Buscemi* (Nagoya U)

*www.quantumquia.com

KAIST-Nagoya GENKO Workshop, Daejeong, 21 February 2023

collaborators

- Clive Aw (CQT@NUS)
- Arthur Parzygnat (Nagoya)
- Dominik Šafránek (IBS)
- Valerio Scarani (CQT@NUS)
- Joseph Schindler (UAB)

The Observational Entropy Appreciation Club

(www.observationalentropy.com)

von Neumann's entropy

For $\varrho = \sum_{x=1}^{d} \lambda_x |\varphi_x\rangle \langle \varphi_x | d$ -dimensional density matrix ($\lambda_x \ge 0$, $\sum_x \lambda_x = 1$),

$$S(\varrho) := -\operatorname{Tr}[\varrho \log \varrho] = -\sum_{x=1}^{d} \lambda_x \log \lambda_x$$

with the convention $0 \log 0 := 0$.

Unfortunately though:

"The expressions for entropy given by the author [previously] are not applicable here in the way they were intended, as they were computed from the perspective of an observer who can carry out all measurements that are possible in principle—i.e., regardless of whether they are macroscopic [or not]."

von Neumann, 1929; transl. available in arXiv:1003.2133

3/25

in formula:

Theorem (least uncertainty)

For ρ density matrix, $\mathfrak{onb} = \{ |\phi_i\rangle \}_i$ orthonormal basis, and $p_i = \langle \phi_i | \rho | \phi_i \rangle$,

$$S(\varrho) = \min_{\mathfrak{onb}} \left[-\sum_i p_i \log p_i \right]$$

For a more general result, see [M. Dall'Arno and F.B., IEEE TIT, 65(4), 2018].

5/25

"Although our entropy expression, as we saw, is completely analogous to the classical entropy, it is still surprising that it is invariant in the normal [Hamiltonian] evolution in time of the system, and only increases with measurements—in the classical theory (where the measurements in general played no role) it increased as a rule even with the ordinary mechanical evolution in time of the system. It is therefore necessary to clear up this apparently paradoxical situation."

von Neumann, book (Math. Found. QM), 1932 (transl. 1955)



invariance of von Neumann entropy

Instead,

Theorem

For any unitary operator U,

$$S(\varrho) = S(U\varrho U^{\dagger}) ,$$

for all density matrices ϱ .

 \implies the entropy increasing during a free expansion cannot be the von Neumann entropy

von Neumann's insight (inspired by Szilard's)

"For a classical observer, who knows all coordinates and momenta, the entropy is constant. [...]

The time variations of the entropy are then based on the fact that the observer does not know everything—that he cannot find out (measure) everything which is measurable in principle."

von Neumann, 1932 (transl. 1955)

von Neumann's proposal: macroscopic entropy

For

- ϱ density matrix,
- $\mathfrak{P} = {\{\Pi_i\}_i \text{ orthogonal resolution of identity,}}$
- $p_i = \operatorname{Tr}[\varrho \ \Pi_i]$,
- $\Omega_i := \operatorname{Tr}[\Pi_i]$,

$$S_{\mathfrak{P}}(\varrho) := -\sum_{i} p_i \log \frac{p_i}{\Omega_i}$$

modern generalization: observational entropy For

- ϱ density matrix,
- $\mathbf{P} = \{P_i\}_i \text{ POVM (i.e., } P_i \ge 0, \sum_i P_i = 1),$
- $p_i = \operatorname{Tr}[\varrho \ P_i]$,
- $V_i := \operatorname{Tr}[P_i]$,

$$S_{\mathbf{P}}(\varrho) := -\sum_{i} p_i \log \frac{p_i}{V_i}$$

References:

- D. Šafránek, J.M. Deutsch, A. Aguirre. *Phys. Rev. A* 99, 012103 (2019)
- D. Šafránek, A. Aguirre, J. Schindler, J. M. Deutsch. Found. Phys. 51, 101 (2021)

"observational" = "of the observer"

- von Neumann defines a macro-observer as a collection of simultaneously measurable quantities {Q₁, Q₂, ..., Q_n, ...}, where Q_n = {Q_{x|n}}_x are POVMs
- \implies there exists one "mother" POVM $\mathbf{P} = \{P_i\}_i$ and a stochastic processing (i.e., cond. prob.) μ such that

$$Q_{x|n} = \sum_{i} \mu(x|n,i) P_i , \quad \forall x, n$$

 hence, a "macro-observer" is just a POVM, from which all macroscopic measurements (i.e., coarse-grainings) can be simultaneously inferred by stochastic post-processing

Mathematical properties

from arXiv:2209.03803

Umegaki's relative entropy

Definition

For density matrices ϱ, σ ,

 $D(\varrho \| \sigma) := \begin{cases} \operatorname{Tr}[\varrho(\log \varrho - \log \sigma)] \ , & \text{if } \operatorname{supp} \varrho \subseteq \operatorname{supp} \sigma \ , \\ +\infty \ , & \text{otherwise} \end{cases}$

Useful properties:

- $D(A||B) \ge 0$
- $S(\varrho) = \log d D(\varrho \| u)$ where $u := d^{-1} \mathbb{1}$
- monotonicity: $D(\varrho \| \sigma) \ge D(\mathcal{E}(\varrho) \| \mathcal{E}(\sigma))$ for all channels (i.e., CPTP linear maps) \mathcal{E} and all states ϱ, σ

a bound on the observational entropy

Theorem

For any state ϱ and any POVM $\mathbf{P} = \{P_i\}_i$

 $S(\varrho) \leq S_{\mathbf{P}}(\varrho)$.

Proof.

Given a POVM **P**, by defining the corresponding CPTP linear map $\mathcal{P}(\bullet) := \sum_{i} \operatorname{Tr}[P_i \bullet] |i\rangle\langle i|$, we have $(u = d^{-1}\mathbb{1})$

$$S_{\mathbf{P}}(\varrho) - S(\varrho) = D(\varrho || u) - D(\mathcal{P}(\varrho) || \mathcal{P}(u)) ,$$

which is non-negative due to the monotonicity property of the Umegaki quantum relative entropy.

Petz's theorem

In general, $D(\varrho \| \sigma) \ge D(\mathcal{E}(\varrho) \| \mathcal{E}(\sigma))$. Question: for which triples $(\varrho, \sigma, \mathcal{E})$ does the equality $D(\varrho \| \sigma) = D(\mathcal{E}(\varrho) \| \mathcal{E}(\sigma))$ hold?

Petz (1986,1988)

Answer: if and only if the "transpose channel", i.e.,

$$\widetilde{\mathcal{E}}_{\sigma}(\bullet) := \sqrt{\sigma} \mathcal{E}^{\dagger} \left[\frac{1}{\sqrt{\mathcal{E}(\sigma)}} \bullet \frac{1}{\sqrt{\mathcal{E}(\sigma)}} \right] \sqrt{\sigma}$$

satisfies $\widetilde{\mathcal{E}}_{\sigma} \circ \mathcal{E}(\varrho) = \varrho$. (The other equality $\widetilde{\mathcal{E}}_{\sigma} \circ \mathcal{E}(\sigma) = \sigma$ is satisfied by construction.)

Remark. Notice that $\widetilde{\mathcal{E}}_{\sigma}$ in general is *not* the linear inverse of \mathcal{E} ; rather, it is related with the idea of *statistical retrodiction* (more on this later).

15/25

consequences for observational entropy

Theorem

$$S(\varrho) = S_{\mathbf{P}}(\varrho) \iff \varrho = \sum_{i} \operatorname{Tr}[\varrho \ P_i] \frac{P_i}{V_i}.$$

We call such a state fully observable or macroscopic (w.r.t. observer **P**).

Proof.

This is a direct consequence of Petz's transpose map theorem: $S_{\mathbf{P}}(\varrho) - S(\varrho) = D(\varrho || u) - D(\mathcal{P}(\varrho) || \mathcal{P}(u)) = 0$ if and only if $\varrho = \widetilde{\mathcal{P}}_u(\mathcal{P}(\varrho))$. By direct inspection, $\widetilde{\mathcal{P}}_u(\mathcal{P}(\varrho)) = \sum_i \operatorname{Tr}[\varrho \ P_i] P_i/V_i$.

Remark. For any observer (i.e., POVM) **P**, we denote the subspace of macroscopic states by $\mathfrak{M}(\mathbf{P})$.

```
17/25
```

the resolution of the paradox

- start at $t = t_0$ from a macrostate $\varrho^{t_0} \in \mathfrak{M}(\mathbf{P})$
- the system undergoes unitary evolution, i.e., $\varrho^{t_0} \mapsto \varrho^{t_1} = U \varrho^{t_0} U^{\dagger}$; then,

$$\begin{split} S_{\mathbf{P}}(\varrho^{t_1}) &= -\sum_i \operatorname{Tr} \left[P_i \left(U \varrho^{t_0} U^{\dagger} \right) \right] \log \frac{\operatorname{Tr} \left[P_i \left(U \varrho^{t_0} U^{\dagger} \right) \right]}{\operatorname{Tr} \left[P_i \right]} \\ &= -\sum_i \operatorname{Tr} \left[\left(U^{\dagger} P_i U \right) \varrho^{t_0} \right] \log \frac{\operatorname{Tr} \left[\left(U^{\dagger} P_i U \right) \varrho^{t_0} \right]}{\operatorname{Tr} \left[U^{\dagger} P_i U \right]} \\ &= S_{U^{\dagger} \mathbf{P} U}(\varrho^{t_0}) \\ &\geq S_{\mathbf{P}}(\varrho^{t_0}) = S(\varrho^{t_0}) = S(\varrho^{t_1}) \end{split}$$

• summarizing: in general, $S_{\mathbf{P}}(\varrho^{t_1}) \ge S_{\mathbf{P}}(\varrho^{t_0})$

- with equality if and only if $\varrho^{t_1} \in \mathfrak{M}(\mathbf{P})$
- in words: the observational entropy of an isolated macroscopic state can (in fact, will generically) increase, and it remains constant if and only if the state remains macroscopic

Generalization to finite differences

triangle equality for observational entropy

Theorem

Given a d-dimensional system, a density matrix ρ with diagonalization $\{\lambda_x, |\varphi_x\rangle\}_{x=1}^d$, a unitary operator U, and a POVM $\mathbf{P} = \{P_i\}_i$, let us define two joint probability distributions:

$$P_F(x,i) := \lambda_x \underbrace{\operatorname{Tr}\left[U|\varphi_x\rangle\!\langle\varphi_x|U^{\dagger} P_i\right]}_{P_F(i|x)},$$

$$P_R(x,i) := p_i \underbrace{\operatorname{Tr}\left[|\varphi_x\rangle\!\langle\varphi_x| \; \frac{U^{\dagger} P_i U}{V_i}\right]}_{P_R(x|i)}$$

Then,
$$S_{\mathbf{P}}(U\varrho U^{\dagger}) - S(\varrho) = D(P_F || P_R).$$



interpretation: prediction and retrodiction

in the language of Jeffrey's "probability kinematics"

- P_F corresponds to the prediction $\lambda \xrightarrow{\pi} \bullet$: the inference about i
- P_R corresponds to the retrodiction ^π/_u p: the inference about x that a completely uninformed Bayesian agent would do, if given information about i in the form of the probability distribution p.

The equality $S_{\mathbf{P}}(U\varrho U^{\dagger}) = S(\varrho)$ occurs if and only if predictor and retrodictor agree.

Watanabe's contention



"The phenomenological onewayness of temporal developments in physics is due to irretrodictability, and not due to irreversibility." Satosi Watanabe (1965)

The second law of thermodynamics is not about the "arrow of time", but about the arrow of inference.

- F.B. and V. Scarani, Fluctuation relations from Bayesian retrodiction, PRE (2021)
- C.C. Aw, F.B., and V. Scarani, Fluctuation theorems with retrodiction rather than reverse processes, AVS Quantum Science (2021)

a double bound for quantum retrodiction

Can we say something about the quantum process itself, i.e., independently of any particular diagonalization of ρ ?

Theorem

Given a density matrix ϱ , a unitary operator U, and a POVM $\mathbf{P} = \{P_i\}_i$,

$$T\ln(d-1) + h(T) \ge S_{\mathsf{P}}(U\varrho U^{\dagger}) - S(\varrho) \ge D(\varrho \| \widetilde{\varrho}_{\mathbf{r}}) ,$$

where $T := \frac{1}{2} \| \varrho - \widetilde{\varrho}_r \|_1$, $h(x) := -x \ln x - (1-x) \ln(1-x)$, and $\widetilde{\varrho}_r := \sum_i \operatorname{Tr} \left[U \varrho U^{\dagger} P_i \right] \frac{U^{\dagger} P_i U}{V_i}$ is the state retrodicted by the later observer.

Remark. The above bound is genuinely quantum, i.e., $[\varrho, \tilde{\varrho}_r] \neq 0$.

23/25

Conclusions

take-home messages

When the use of von Neumann entropy in thermodynamics is problematic, try consider observational entropy (OE) instead, because:

- OE has a fully operational/inferential definition
- OE describes thermodynamic scenarios avoiding interpretational paradoxes
- OE fits nicely within recent developments in quantum mathematical statistics (e.g., approximate Petz recovery)
- OE suggests a natural candidate for quantum retrodiction

The End: Thank You!